Seroprevalence of Syphilis, CMV and Rubella Infection in Waste-Blood Samples

Nyein Ko Hein, Than Naing Oo, Zaw Min Tun, Aung Zay Hein, Tin Moe Mya, Thein Zaw & Tin Maung Hlaing Defence Services Medical Research Centre, Nay Pyi Taw, Myanmar

dr.nyeinkohein@gmail.com


❖ Waste-blood samples after testing for Non-communicable diseases

MYANMAR


- From 4 hospitals in Nay Pyi Taw, Myanmar including 1 OG hospital.
- * Test: LFT, Lipid Profile, U&E, Cardiac Enzymes, RA, SLE, Hormones, T&DP etc.
- * RPR carbon kit for Syphilis, ELISA for IgG of CMV and Rubella


Number of Examined	Syphilis (Positive) No. (%)	CMV (IgG) (Positive) No. (%)	Rubella (lgG) (Positive) No. (%)	Syphilis + CMV + Rubella (Positive) No. (%)	Syphilis+ CMV (Positive) No. (%)	CMV + Rubella (Positive) No. (%)
500	3 (0.6)	362 (72.4)	350 (70.0)	2 (0.4)	1 (0.2)	285 (57.0)











- Screen Active Infection and Treat
- ❖ Prevention of Syphilis and Rubella Infection

THANK YOU at See you at Poster 26