

GENETIC VARIATIONS IN *AEROMONAS HYDROPHILA* ISOLATES FROM CLINICAL AND ENVIRONMENTAL SOURCES IN THAILAND

S Korbsrisate¹, S Trakulsomboon², S Damnin¹, J Gatedee³, OA Ratchtrachenchai⁴,
A Leelaporn⁵, B Eampokalap⁶, C Gongviseisoog⁶ and S Suparak¹

¹Department of Immunology, ²Department of Medicine, ⁵Department of Microbiology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok; ³Faculty of Medical Technology, Rangsit University, Pathum Thani; ⁴Department of Medical Sciences, National Institute of Health, Nonthaburi; ⁶Microbiology Laboratory, Bamrasnaradura Hospital, Nonthaburi, Thailand

Abstract. *Aeromonas hydrophila*, a widely distributed human pathogen causing a variety of diseases, can be isolated from clinical and environmental sources. Analysis in Thailand of 110 isolates of *Aeromonas hydrophila* by randomly amplified polymorphic DNA-PCR (RAPD-PCR) revealed one specific RAPD pattern group (G) that was associated only with strains from environmental sources. Cytotoxic activity, adhesion to epithelial cells and exoenzyme secretions of *A. hydrophila* were also investigated. A comparison of isolates with pattern group G with a set of isolates derived from human blood showed low induction of cytotoxicity from those with RAPD pattern group G suggesting low virulence of these strains.

Correspondence: S Korbsrisate, Department of Immunology, Faculty of Medicine Siriraj Hospital, Mahidol University, 2 Pran Nok Road, Bangkok 10700, Thailand.
E-mail: grsks@mahidol.ac.th