

## REVIEW

# SEROLOGICAL AND MOLECULAR TOOLS TO DETECT NEUROLOGIC PARASITIC ZOOSES IN RURAL CAMEROON

Agathe Nkouawa<sup>1,2,a</sup>, Yasuhito Sako<sup>1</sup>, Roger Moyou-Somo<sup>2,3</sup> and Akira Ito<sup>1,a</sup>

<sup>1</sup>Department of Parasitology, Asahikawa Medical University, Asahikawa, Hokkaido, Japan; <sup>2</sup>Medical Research Center, Institute of Medical Research and Medicinal Plants Studies, Ministry of Scientific Research and Innovation, Yaoundé; <sup>3</sup>Department of Parasitology and Infectious Diseases, Faculty of Medicine and Biomedical Sciences, University of Yaounde I, Yaoundé, Cameroon

**Abstract.** Parasitic helminthiasis, such as toxocariasis, cysticercosis and paragonimiasis are a public health threat, since they can affect the brain leading to neurological disorders. Epilepsy and paragonimiasis are common in southwestern Cameroon. We reviewed the literature for studies using antigens to diagnose toxocariasis, cysticercosis, and paragonimiasis. Serology revealed that 61 (36.3%), 26 (15.5%) and 2 (1.2%) of 168 persons examined [78 males (15.2 ± 8.2 years old), 90 females (12.9 ± 5.9 years old), 143 persons < 20 years old] had antibody responses to toxocariasis, paragonimiasis and cysticercosis, respectively. Of the 14 people with epilepsy, 5 were seropositive for *Toxocara* antigens and 1 was positive for both *Toxocara* and *Paragonimus* antigens. Two children were serologically confirmed to have cysticercosis. Serologic screening for cysticercosis may be feasible to detect asymptomatic cysticercosis in children in endemic areas leading to early treatment. The causative *Paragonimus* species was confirmed to be *P. africanus* by molecular sequencing. Education, screening and confirmation test for these diseases may be needed for control in Cameroon.

**Keywords:** epilepsy, parasitic zoonoses, toxocariasis, paragonimiasis, cysticercosis, immuno- and molecular-diagnosis, Cameroon

## INTRODUCTION

Parasitic zoonotic infections have a potentially serious impact on human

---

Correspondence: Dr Agathe Nkouawa, Medical Research Center, Institute of Medical Research and Medicinal Plants Studies, Ministry of Scientific Research and Innovation, PO Box 6163, Yaoundé, Cameroon.

E-mail: ankouawa@yahoo.fr

<sup>a</sup>These authors contributed equally.

health and economy. Paragonimiasis, cysticercosis and toxocariasis represent a public health threat in developing countries, since they can affect various tissues and organs including the brain leading to neurological disorders, such as epilepsy (Nicoletti *et al*, 2002; Ito *et al*, 2006b; Garcia and Modi, 2008). Cysticercosis has been considered to be a major cause of the late-onset epilepsy in developing countries (Zoli *et al*, 2003; Montano *et al*,

2005; Villaran *et al*, 2009). Other helminthic diseases such as toxocariasis, paragonimiasis, and onchocerciasis and protozoan diseases, such as malaria and toxoplasmosis, can also cause neurological disorders and idiopathic epilepsy (Bachli *et al*, 2004; Akyol *et al*, 2007; Garcia and Modi, 2008; Kaiser *et al*, 2008; Ngoungou and Preux, 2008). Therefore, early detection of these pathogens in humans using immunological and molecular tools is important for early treatment and prevention of the neurological sequelae (Ito, 2002; Ito and Craig, 2003; Margono *et al*, 2003; Ito *et al*, 2006b).

In general, ELISA using crude antigens is suitable for screening and the immunoblot using crude antigens or semi-purified antigens is for confirmation in the diagnosis of these helminthes. With pure antigens the ELISA and immunoblot may have a better specificity. Both these tools might be used to screen and confirm infections in endemic areas (Ito *et al*, 1998, 1999; Sako *et al*, 2000; Sato *et al*, 2003, 2006).

Tombel Health District, Southwest Province, Cameroon is endemic for paragonimiasis and epilepsy is common (Moyou-Somo *et al*, 2003; Moyou-Somo and Tagni-Zukam, 2003). However, there is a paucity of information about the role of parasitic zoonoses among people with neurological disorders in this area. There is little information about cysticercosis in children, although cysticercosis is relatively common in adults and epileptic patients (Nguekam *et al*, 2003; Zoli *et al*, 2003). There is no data regarding the prevalence of toxocariasis in human in Cameroon, although toxocariasis is prevalent in dogs (Komtangi *et al*, 2005). Paragonimiasis is diagnosed by the detection of eggs in the sputum but control of this disease has only been partially successful.

We reviewed the literature regarding the prevalence of toxocariasis, paragonimiasis and cysticercosis, and their association with neurological diseases among children in southwestern Cameroon using immunological techniques.

## TOXOCARIASIS

Toxocariasis is a zoonotic infection of humans caused by dog and cat roundworms, *Toxocara canis* and *Toxocara cati*, respectively. Humans become infected by ingestion of embryonated eggs from contaminated sources. Children often become infected because they put objects in their mouths or eat dirt. This disease can be found in developed and developing countries (Hayashi *et al*, 2005; Fernando *et al*, 2007; Nicoletti *et al*, 2008; Yoshikawa *et al*, 2008; Zarnowska *et al*, 2008). We expect *T. canis* infections are more common than *T. cati* infections due to the defecation habits of dogs, but there is no information about this disease in humans in Cameroon. The prevalence of this disease in dogs is high in Cameroon (Komtangi *et al*, 2005). We expected the prevalence of this infection in children might also be high especially in rural areas where risk factors are common in Cameroon.

Toxocariasis affecting the organs is called visceral larva migrans, and the disease restricted to the eye and the optic nerve is called ocular larva migrans (Logar *et al*, 2004; Yoshikawa *et al*, 2008). Neurological disorders, especially epilepsy have been associated with toxocariasis (Bachli *et al*, 2004; Moreira-Silva *et al*, 2004; Akyol *et al*, 2007; Nicoletti *et al*, 2008). Epilepsy is common in southwestern Cameroon. Serology using a recombinant antigen of *T. canis* second-stage larvae (Yamasaki *et al*, 2000) was performed on serum samples collected from children in Southwest

Table 1  
Clinical and serological results of individuals investigated.

Clinical/immunological diagnosis of persons with symptoms	Age/gender (male/female)			Total (%) (n = 168)
	0-10 years (31/49)	11-20 years (35/28)	> 21 years (14/11)	
Cough	64 (31/33)	51 (29/22)	20 (12/8)	135 (80.3)
Headache	49 (22/27)	40 (22/18)	17 (7/10)	106 (63.0)
Eye disorder	7 (3/4)	9 (5/4)	14 (6/8)	30 (17.8)
Epilepsy	2 (2/0)	7 (5/2)	5 (3/2)	14 (8.3)
Chest pain	33 (15/18)	30 (15/15)	17 (7/10)	80 (47.6)
Hemoptysis	8 (3/5)	7 (3/4)	3 (1/2)	18 (11.3)
Crab eaten	60 (26/34)	60 (34/26)	17 (11/6)	137 (81.5)
Pork eaten	64 (26/38)	49 (28/21)	22 (12/10)	135 (80.3)
<b>Microscopy for <i>Paragonimus</i> spp eggs<sup>a</sup></b>				
Sputum	9 (2/7)	5 (3/2)	2 (0/2)	16 (9.5)
Feces	0 (0/0)	0 (0/0)	0 (0/0)	0 (0.0)
<b>ELISA and/or immunoblot</b>				
Paragonimiasis	13 (4/9)	9 (5/4)	4 (2/2)	26 (15.5)
Toxocariasis	31 (13/18)	24 (14/10)	6 (4/2)	61 (36.3)
Cysticercosis	0 (0/0)	2 (2/0)	0 (0/0)	2 (1.2)

<sup>a</sup>The eggs were later confirmed to be *P. africanus* using molecular tools.

Cameroon. ELISA results showed that 61 persons (36.3%) exhibited antibody response to this antigen (Table 1) (Nkouawa *et al*, 2010a). It suggests that children are the most exposed people (Hayashi *et al*, 2005; Fernando *et al*, 2007; Zarnowska *et al*, 2008; Sviben *et al*, 2009). Among the 14 persons with epilepsy, 5 were seropositive for specific *Toxocara* antigen (Fig 1). More epidemiological data of toxocariasis are needed to determine its association with epilepsy in this area, and in other regions in Cameroon to determine the distribution of this disease.

### PARAGONIMIASIS

Paragonimiasis is a parasitic disease of humans and other mammals caused by *Paragonimus* spp (lung flukes). Humans are infected after ingestion of raw or un-

dercooked fresh water crabs harboring the metacercariae. Pulmonary paragonimiasis is the most common, but cerebral paragonimiasis is not rare. Since both paragonimiasis and epilepsy are highly prevalent in the Southwest Province of Cameroon, it is necessary to investigate the correlation between epilepsy and paragonimiasis in this area. Microscopic examination of eggs in feces is generally the most reliable direct method. However, *Paragonimus* eggs were mostly found from sputa but not from fecal samples collected in Southwest Cameroon (Nkouawa *et al*, 2009a). This might be due to the people's attitude to vomit sputa rather than swallowing them. Therefore, the diagnosis of paragonimiasis in Cameroon by detecting *Paragonimus* eggs from feces lacks sensitivity (Ripert *et al*, 1992; Ollivier *et al*, 1995; Moyou-Somo *et al*, 2003; Moyou-Somo




Fig 1–ELISA results modified from Nkouawa *et al* (2010a) for paragonimiasis (Pa), cysticercosis (Ts) and toxocariasis (Tc) from 208 persons [168 and 20 (Nc) with and without symptoms from Cameroon, respectively and 20 from Japan (Nj)]. ○, samples with *Paragonimus* eggs in the sputum. Serology could detect more than all cases with eggs in the sputum and was expected to be more sensitive for detection of paragonimiasis including immature adult stage (Nkouawa *et al*, 2009a). ◇, positive against the recombinant antigen 100% specific to cysticercosis by immunoblot (Sako *et al*, 2000). △, patients with history of seizures were positive to *P. africanus* somatic antigen and to recombinant antigen of *T. canis* second-stage larvae. The broken line denotes the respective cut-off value for each disease.

and Tagni-Zukam, 2003). Furthermore, it is not possible to detect eggs during pre-patent period, in single infections, or extra-pulmonary paragonimiasis. Serological tests, which have been confirmed to be useful for detection of paragonimiasis in Asia and Latin America (Ikeda *et al*, 1996; Kong *et al*, 1998; Blair *et al*, 2007), are more sensitive than detection of eggs in the sputum for diagnosis of paragonimiasis in Cameroon (Nkouawa *et al*, 2009a). A positive antibody responses to crude antigens of *P. africanus* adult worms were observed in 14.8% ( $n = 25$ ) and 15.5% ( $n = 26$ ) of 168 sera collected from persons living in Southwest Cameroon by enzyme-linked immunosorbent assay (ELISA) and immunoblot analysis, respectively (Table 1). Sera from other parasitic infections (schistosomiasis haematobium, schistosomiasis japonicum, fascioliasis, clonorchiasis, cysticercosis and spargano-

sis) showed weak cross-reaction to crude antigens (Nkouawa *et al*, 2009a). Immunodominant bands of 33- and 35-kDa of *P. africanus* antigens recognized by most patients' sera were similar to those of *P. westermani* adults exhibiting strong reactions with IgG4 (Kong *et al*, 1998). Crude antigens of *Paragonimus* spp are useful for detection of paragonimiasis, since these antigens appear to be genus specific (Kong *et al*, 1998). Sera from persons infected with *P. africanus* showed strong antibody response to the Japanese *P. westermani* and *P. miyazakii* adult antigens, and *vice versa* sera from persons infected with the Japanese *P. westermani* and *P. miyazakii* strongly cross-reacted with the antigens of *P. africanus* adult (Nkouawa *et al*, 2009a). This indicates that differentiation of *Paragonimus* species cannot be achieved serologically. Since at least ten *Paragonimus* species have been reported to infect

humans in endemic areas in Asia (Sripa *et al*, 2010), antigens from any species are expected to provide reliable results.

In Cameroon, immunological tests detected more paragonimiasis cases (15.4%) than microscopic examination of eggs in the sputum (3.0-9.6%) (Ripert *et al*, 1992; Ollivier *et al*, 1995; Moyou-Somo *et al*, 2003; Moyou-Somo and Tagni-Zukam, 2003). Therefore, we adopted immunological tests for the surveillance of this disease. Although the number of cases has decreased in adult populations in endemic areas in Southwest Cameroon (Moyou-Somo *et al*, 2003; Moyou-Somo and Tagni-Zukam, 2003), our data have demonstrated that children, especially those < 10 years old are still highly infected with *P. africanus* due to their food eating habits and poor hygiene and sanitation (Table 1). Therefore, control measures including mass screening by immunological method and education of children are an urgent task to prevent this disease. Out of 14 epileptic patients, only one showed positive antibody responses to antigens specific to paragonimiasis (Fig 1), this suggests that paragonimiasis might not be the major cause of epilepsy in this area.

Since four species, *P. africanus*, *P. uterobilateralis*, *P. westermani*-like and *Euparagonimus* sp, are distributed in Cameroon (Voelker and Sachs, 1977; Cabaret *et al*, 1999), it is necessary to confirm the parasite species causing paragonimiasis by molecular tools. Adult worms recovered from the lungs of a cat after experimental infection with metacercariae collected from crabs in Southwest Cameroon were identified as *P. africanus* by sequence data from the internal transcribed spacer 2 (ITS2) region and cytochrome *c* oxidase subunit 1 (cox1) genes (Nkouawa *et al*, 2009a). Therefore, molecular sequencing can be used to detect *P. africanus*,

the causative agent of paragonimiasis in Southwest Cameroon. PCR using specific primer designed from *P. africanus* ITS2 and cox1 sequences was also used to detect target DNA from fecal samples. It is more sensitive than microscopic examination of eggs in feces (Nkouawa *et al*, 2009a).

Detection of target DNA in the sputum provides more sensitive results, especially to identify the *Paragonimus* species, since eggs are often found in cases of pulmonary paragonimiasis. This method cannot be applied for extra-pulmonary cases or cerebral paragonimiasis.

## CYSTICERCOSIS

Cysticercosis caused by *T. solium* larvae is one of the most serious diseases in many developing countries (Schantz *et al*, 1998; Murrell, 2005; Ito *et al*, 2006a). Previous studies have focused on adult population and pigs in the western and northwestern regions of Cameroon (Pouedet *et al*, 2002; Vondou *et al*, 2002; Nguekam *et al*, 2003; Shey-Njila *et al*, 2003; Zoli *et al*, 2003), and on swine cysticercosis in the North Province (Assana *et al*, 2010). There is little or no information from other regions, especially where epilepsy is relatively high and where the risk factor of this disease such as pig farming with free access of pigs to human feces remains. For human cysticercosis, detection of antibody responses has been combined with clinical criteria and/or imaging figures (Ito *et al*, 2006c; Willingham *et al*, 2010). As the CT or MRI is too expensive and is not affordable to many people in developing countries, immunodiagnosis using specific antigens is the method of choice for screening of cysticercosis due to its simplicity and reliability.

Among several characterized specific antigens for *T. solium*, the glycoproteins



(GPs) have been widely used for serodiagnosis of cysticercosis (Ito *et al*, 1998; Sako *et al*, 2000; Wandra *et al*, 2003; Ito *et al*, 2006c; Li *et al*, 2006; Sato *et al*, 2006; Wandra *et al*, 2006). We applied GPs and recombinant antigen (Sako *et al*, 2000; Li *et al*, 2006; Sato *et al*, 2006; Anantaphruti *et al*, 2010) for diagnosis of cysticercosis in serum samples collected in Southwest Cameroon (Nkouawa *et al*, 2010a). More than half of epileptic adult patients in West and North West provinces of Cameroon exhibited strong antibody responses. It suggested that cysticercosis was the major causative agent of the late-onset epilepsy in this area (Zoli *et al*, 2003). However, out of 168 samples including 143 persons < 20 years old, collected from Southwest Cameroon, 1 girl aged 4 years old and 2 boys aged 11 and 13 years old without symptoms or any history of seizures exhibited antibody responses against GPs by ELISA. Immunoblot using the same GPs and recombinant antigens of *T. solium* revealed that the 2 boys were seropositive. Therefore, these 2 children are asymptomatic cysticercosis (Nkouawa *et al*, 2010a). As it has been shown that cysticercus of *T. solium* can survive for many years in patient's body (Yanagida *et al*, 2010), it is necessary to follow up these two children.

The follow-up of asymptomatic healthy but serologically positive people for cysticercosis in Papua, Indonesia, revealed that most of them had detectable subcutaneous nodules (Wandra *et al*, 2003; Ito *et al*, 2004). However, subcutaneous cysticercosis is not common in Africa as in Asia (Ito *et al*, 2003). Therefore, follow-up antibody titers may be another indicator to confirm active cysticercosis in Africa. Serology using highly specific antigens is useful for detection of asymptomatic cysticercosis and symptomatic

cases in endemic areas.

## TAENIASIS

To achieve the control and prevention of cysticercosis, the detection of cysticercosis in humans, pigs, and dogs (Ito *et al*, 2002) together with the detection of taeniasis carriers are required in endemic areas. Although cysticercosis is still a serious problem in some areas in Asia, Africa and Latin America, the disease has been well controlled (Ito *et al*, 2005; Garcia *et al*, 2007; Wandra *et al*, 2006). Nonetheless, the control in Cameroon is mainly focused on the detection of cysticerci in humans and pigs, which are the end stage of the parasite in the intermediate hosts. There is few information on adult tapeworms and the risk factor for cysticercosis in pigs and humans in many endemic areas. A previous study reported that *T. solium* taeniasis was diagnosed by microscopic examination of *Taenia* egg in fecal samples collected from people in endemic areas of West Cameroon (Vondou *et al*, 2002).

The microscopic examination of *Taenia* eggs is not sensitive and eggs of all *Taenia* species are morphologically identical. More reliable molecular tools such as multiplex PCR and loop-mediated isothermal amplification (LAMP) have been introduced for identification of eggs, cysticerci, adult worms and for feces without eggs or proglottids (Nkouawa *et al*, 2009b; Yamasaki *et al*, 2004). LAMP is more sensitive to detect taeniasis than multiplex PCR (Nkouawa *et al*, 2010b) in endemic areas in Cameroon. For future studies in endemic areas in Cameroon and other countries, the detection of adult tapeworms as a screening of taeniasis carriers of *T. solium* using both copro-ELISA (Guezala *et al*, 2009) and copro-DNA methods (Nkouawa *et al*, 2009b, 2010b; Nakao *et al*, 2010) is necessary.

## CONCLUSION

In this review, highly specific recombinant antigens for both toxocariasis and cysticercosis, and crude antigens of *P. africanus* showing no or little cross-reaction with other helminth infections were used. Toxocariasis (36.3%), paragonimiasis (15.5%) and cysticercosis (1.2%) have been confirmed among surveyed persons. Among the seropositive persons, 11 were found to have dual infections of both *Toxocara* and *Paragonimus*. Of 14 epilepsy cases, 5 were seropositive for toxocariasis and one was simultaneously positive for paragonimiasis. This demonstrated that 1) paragonimiasis among children still remains a serious health problem in this endemic area, 2) asymptomatic cysticercosis cases were detected in 2 children, and 3) toxocariasis was serologically confirmed for the first time in Cameroon with relatively high prevalence. The serological tools are highly useful for screening of these diseases having great consequences on the brain. Therefore, early diagnosis at the early stage is one of the key strategies to control and prevent the spread of these diseases in people, especially in children in endemic areas. For future strategies, detail epidemiological studies to screen young and adult populations for taeniasis/cysticercosis, and pigs for cysticercosis and also other helminthic infections are necessary to establish the correlation between epilepsy and parasitic diseases for the early treatment and prevention of these diseases in this area and other regions in Cameroon and any other countries.

## ACKNOWLEDGEMENTS

This work was supported in part by the Special Coordination Funds for Promoting Science and Technology from the

Ministry of Education, Culture, Sports, Science and Technology, Japan (2010-2012); the International Collaboration Research Fund from the Japan Society for the Promotion of Science (JSPS) (17256002, 21256003) and JSPS-Asia/Africa Science Platform Fund (2006-2011) to AI.

## REFERENCES

- Akyol A, Bicerol B, Ertug S, Ertabaklar H, Kiylioglu N. Epilepsy and seropositivity rates of *Toxocara canis* and *Toxoplasma gondii*. *Seizure* 2007; 16: 233-7.
- Anantaphruti MT, Okamoto M, Yoonuan T, et al. Molecular and serological survey on taeniasis and cysticercosis in Kanchanaburi Province, Thailand. *Parasitol Int* 2010; 59: 326-30.
- Assana E, Amadou F, Thys E, et al. Pig-farming systems and porcine cysticercosis in the north of Cameroon. *J Helminthol* 2010; 25: 1-6.
- Bachli H, Minet JC, Gratzl O. Cerebral toxocariasis: a possible cause of epileptic seizure in children. *Childs Nerv Syst* 2004; 20: 468-72.
- Blair D, Agatsuma T, Wang W. Paragonimiasis. Fish- and plant-borne parasites. Chapter 3. In: Murrell KD, Fried B, eds. *World class parasites*. New York: Springer, 2007; 11: 117-50.
- Cabaret J, Bayssade-Dufour C, Tami G, Albaret JL. Identification of African Paragonimidae by multivariate analysis of the eggs. *Acta Trop* 1999; 72: 79-89.
- Fernando SD, Wickramasinghe VP, Kapilanda GM, Devasurendra RL, Amarasooriya JD, Dayaratne HG. Epidemiological aspects and risk factors of toxocariasis in a pediatric population in Sri Lanka. *Southeast Asian J Trop Med Public Health* 2007; 38: 983-90.
- Garcia HH, Gonzalez AE, Del Brutto OH, et al. Strategies for the elimination of taeniasis/cysticercosis. *J Neurol Sci* 2007; 262: 153-7.
- Garcia HH, Modi M. Helminthic parasites and

- seizures. *Epilepsia* 2008; 49 (suppl 6): 25-32.
- Guezala MC, Rodriguez S, Zamora H, *et al.* Development of a species-specific coproantigen ELISA for human *Taenia solium* taeniasis. *Am J Trop Med Hyg* 2009; 81: 433-7.
- Hayashi E, Tuda J, Imada M, Akao N, Fujita K. The high prevalence of asymptomatic *Toxocara* infection among schoolchildren in Manado, Indonesia. *Southeast Asian J Trop Med Public Health* 2005; 36: 1399-406.
- Ikeda T, Oikawa Y, Nishiyama T. Enzyme-linked immunosorbent assay using cysteine proteinase antigens for immunodiagnosis of human paragonimiasis. *Am J Trop Med Hyg* 1996; 55: 435-7.
- Ito A. Serologic and molecular diagnosis of zoonotic larval cestode infections. *Parasitol Int* 2002; 51: 221-35.
- Ito A, Craig PS. Immunodiagnostic and molecular approaches for the detection of taeniid cestode infections. *Trends Parasitol* 2003; 19: 377-81.
- Ito A, Craig PS, Schantz PM. Taeniasis/cysticercosis and echinococcosis with focus on Asia and the Pacific. *Parasitol Int* 2006a; 55 (suppl): 1-312.
- Ito A, Nakao M, Wandra T, *et al.* Taeniasis and cysticercosis in Asia and the Pacific: present state of knowledge and perspectives. *Southeast Asian J Trop Med Public Health* 2005; 36 (suppl 4): 123-30.
- Ito A, Plancarte A, Ma L, *et al.* Novel antigens for neurocysticercosis: simple method for preparation and evaluation for serodiagnosis. *Am J Trop Med Hyg* 1998; 59: 291-4.
- Ito A, Plancarte A, Nakao M, *et al.* ELISA and immunoblot using purified glycoproteins for serodiagnosis of cysticercosis in pigs naturally infected with *Taenia solium*. *J Helminthol* 1999; 73: 363-5.
- Ito A, Putra MI, Subahar R, *et al.* Dogs as alternative intermediate hosts of *Taenia solium* in Papua (Irian Jaya), Indonesia confirmed by highly specific ELISA and immunoblot using native and recombinant antigens and mitochondrial DNA analysis. *J Helminthol* 2002; 76: 311-4.
- Ito A, Takayanagui OM, Sako Y, *et al.* Neurocysticercosis: clinical manifestation, neuroimaging, serology and molecular confirmation of histopathologic specimens. *Southeast Asian J Trop Med Public Health* 2006b; 37(suppl 3): 74-81.
- Ito A, Wandra T, Sato MO, *et al.* Towards the international collaboration for detection, surveillance and control of taeniasis/ cysticercosis and echinococcosis in Asia and the Pacific. *Southeast Asian J Trop Med Public Health* 2006c; 37(suppl 3): 82-90.
- Ito A, Wandra T, Yamasaki H, *et al.* Cysticercosis/taeniasis in Asia and the Pacific. *Vector Borne Zoonotic Dis* 2004; 4: 95-107.
- Ito A, Yamasaki H, Nakao M, *et al.* Multiple genotypes of *Taenia solium*—ramifications for diagnosis, treatment and control. *Acta Trop* 2003; 87: 95-101.
- Kaiser C, Pion S, Preux PM, Kipp W, Dozie I, Boussinesq M. Onchocerciasis, cysticercosis, and epilepsy. *Am J Trop Med Hyg* 2008; 79: 643-4; author reply 644-5.
- Komtangi MC, Mpoame M, Payne VK, Ngufor MN. Prevalence of gastrointestinal helminths of dogs in Dschang, Cameroon. *J Cameroon Acad Sci* 2005; 5: 11-4.
- Kong Y, Ito A, Yang HJ, *et al.* Immunoglobulin G (IgG) subclass and IgE responses in human paragonimiasis caused by three different species. *Clin Diagn Lab Immunol* 1998; 5: 474-8.
- Li T, Craig PS, Ito A, *et al.* Taeniasis/cysticercosis in a Tibetan population in Sichuan Province, China. *Acta Trop* 2006; 100: 223-31.
- Logar J, Soba B, Kraut A, Stirn-Kranjc B. Seroprevalence of *Toxocara* antibodies among patients suspected of ocular toxocariasis in Slovenia. *Korean J Parasitol* 2004; 42: 137-40.
- Margono SS, Ito A, Sato MO, *et al.* *Taenia solium* taeniasis/cysticercosis in Papua, Indonesia in 2001: detection of human worm carriers. *J Helminthol* 2003; 77: 39-42.
- Montano SM, Villaran MV, Ylquimiche L, *et al.* Neurocysticercosis: association between



- seizures, serology, and brain CT in rural Peru. *Neurology* 2005; 65: 229-33.
- Moreira-Silva SF, Rodrigues MG, Pimenta JL, Gomes CP, Freire LH, Pereira FE. Toxocariasis of the central nervous system: with report of two cases. *Rev Soc Bras Med Trop* 2004; 37: 169-74.
- Moyou-Somo R, Kefie-Arrey C, Dreyfuss G, Dumas M. An epidemiological study of pleuropulmonary paragonimiasis among pupils in the peri-urban zone of Kumba town, Meme Division, Cameroon. *BMC Public Health* 2003; 3: 40.
- Moyou-Somo R, Tagni-Zukam D. Paragonimiasis in Cameroon: clinicoradiologic features and treatment outcome. *Med Trop* 2003; 63: 163-7.
- Murrell KD. WHO/FAO/OIE guidelines for the surveillance, prevention and control of taeniosis/cysticercosis. Paris: OIE, 2005: 1-139.
- Nakao M, Yanagida T, Okamoto M, *et al.* State-of-the-art *Echinococcus* and *Taenia*: phylogenetic taxonomy of human-pathogenic tapeworms and its application to molecular diagnosis. *Infect Genet Evol* 2010; 10: 444-52.
- Ngoungou EB, Preux PM. Cerebral malaria and epilepsy. *Epilepsia* 2008; 49(suppl 6): 19-24.
- Nguekam JP, Zoli AP, Zogo PO, *et al.* A seroepidemiological study of human cysticercosis in West Cameroon. *Trop Med Int Health* 2003; 8: 144-9.
- Nicoletti A, Bartoloni A, Reggio A, *et al.* Epilepsy, cysticercosis, and toxocariasis: a population-based case-control study in rural Bolivia. *Neurology* 2002; 58: 1256-61.
- Nicoletti A, Sofia V, Mantella A, *et al.* Epilepsy and toxocariasis: a case-control study in Italy. *Epilepsia* 2008; 49: 594-9.
- Nkouawa A, Okamoto M, Mabou AK, *et al.* Paragonimiasis in Cameroon: molecular identification, serodiagnosis and clinical manifestations. *Trans R Soc Trop Med Hyg* 2009a; 103: 255-61.
- Nkouawa A, Sako Y, Itoh S, *et al.* Serological studies of neurologic helminthic infections in rural areas of southwest Cameroon: toxocariasis, cysticercosis and paragonimiasis. *PLoS Negl Trop Dis* 2010a; 4: e732.
- Nkouawa A, Sako Y, Li T, *et al.* Evaluation of a loop-mediated isothermal amplification method using fecal specimens for differential detection of *Taenia* species from humans. *J Clin Microbiol* 2010b; 48: 3350-2.
- Nkouawa A, Sako Y, Nakao M, Nakaya K, Ito A. Loop-mediated isothermal amplification method for differentiation and rapid detection of *Taenia* species. *J Clin Microbiol* 2009b; 47: 168-74.
- Ollivier G, Boussinesq M, Albaret JL, *et al.* Epidemiological study of *Paragonimus* sp in south Cameroon. *Bull Soc Pathol Exot* 1995; 88: 164-9.
- Pouedet MS, Zoli AP, Nguekam, *et al.* Epidemiological survey of swine cysticercosis in two rural communities of West-Cameroon. *Vet Parasitol* 2002; 106: 45-54.
- Ripert C, Couprie B, Moyou R, Gaillard F, Appriou M, Tribouley-Duret J. Therapeutic effect of triclabendazole in patients with paragonimiasis in Cameroon: a pilot study. *Trans R Soc Trop Med Hyg* 1992; 86: 417.
- Sako Y, Nakao M, Ikejima T, Piao XZ, Nakaya K, Ito A. Molecular characterization and diagnostic value of *Taenia solium* low-molecular-weight antigen genes. *J Clin Microbiol* 2000; 38: 4439-44.
- Sato MO, Sako Y, Nakao M, Yamasaki H, Nakaya K, Ito A. Evaluation of purified *Taenia solium* glycoproteins and recombinant antigens in the serologic detection of human and swine cysticercosis. *J Infect Dis* 2006; 194: 1783-90.
- Sato MO, Yamasaki H, Sako Y, *et al.* Evaluation of tongue inspection and serology for diagnosis of *Taenia solium* cysticercosis in swine: usefulness of ELISA using purified glycoproteins and recombinant antigen. *Vet Parasitol* 2003; 111: 309-22.
- Schantz PM, Wilkins PP, Tsang VCW. Immigrants, imaging, and immunoblots: the emergence of neurocysticercosis as a significant public health problem. In:

- Scheld WM, Craig WA, Hughes JM, eds. Emerging infections 2. Washington DC: AMS, 1998: 213-42.
- Shey-Njila O, Zoli PA, Awah-Ndukum J, *et al.* Porcine cysticercosis in village pigs of North-West Cameroon. *J Helminthol* 2003; 77: 351-4.
- Sripa B, Kaewkes S, Intapan PM, Maleewong W, Brindley PJ. Food-borne trematodiasis in Southeast Asia: epidemiology, pathology, clinical manifestation and control. *Adv Parasitol* 2010; 72: 305-50.
- Sviben M, Cavlek TV, Missoni EM, Galinovic GM. Seroprevalence of *Toxocara canis* infection among asymptomatic children with eosinophilia in Croatia. *J Helminthol* 2009; 83: 369-71.
- Villaran MV, Montano SM, Gonzalvez G, *et al.* Epilepsy and neurocysticercosis: an incidence study in a Peruvian rural population. *Neuroepidemiology* 2009; 33: 25-31.
- Voelker J, Sachs R. On the distribution of the lung flukes, *Paragonimus africanus* and *P. uterobilateralis*, in the South West Province of Cameroon and in Eastern Nigeria as determined by examination of the intermediate crab hosts for infection with metacercariae. *Tropenmed Parasitol* 1977; 28: 120-33.
- Vondou L, Zoli AP, Nguekam JP, *et al.* *Taenia solium* taeniasis/cysticercosis in the Menoua division (West Cameroon). *Parasite* 2002; 9: 271-4.
- Wandra T, Ito A, Yamasaki H, Suroso T, Margono SS. *Taenia solium* cysticercosis, Irian Jaya, Indonesia. *Emerg Infect Dis* 2003; 9: 884-5.
- Wandra T, Sutisna P, Dharmawan NS, *et al.* High prevalence of *Taenia saginata* taeniasis and status of *Taenia solium* cysticercosis in Bali, Indonesia, 2002-2004. *Trans R Soc Trop Med Hyg* 2006; 100: 346-53.
- Willingham AL 3rd, Wu HW, Conlan J, Satrija F. Combating *Taenia solium* cysticercosis in Southeast Asia an opportunity for improving human health and livestock production. *Adv Parasitol* 2010; 72: 235-66.
- Yamasaki H, Allan JC, Sato MO, *et al.* DNA differential diagnosis of taeniasis and cysticercosis by multiplex PCR. *J Clin Microbiol* 2004; 42: 548-53.
- Yamasaki H, Araki K, Lim PK, *et al.* Development of a highly specific recombinant *Toxocara canis* second-stage larva excretory-secretory antigen for immunodiagnosis of human toxocariasis. *J Clin Microbiol* 2000; 38: 1409-13.
- Yanagida T, Yuzawa I, Joshi DD, *et al.* Neurocysticercosis: assessing where the infection was acquired from. *J Travel Med* 2010; 17: 206-8.
- Yoshikawa M, Nishiofuku M, Moriya K, *et al.* A familial case of visceral toxocariasis due to consumption of raw bovine liver. *Parasitol Int* 2008; 57: 525-9.
- Zarnowska H, Borecka A, Gawor J, Marczyńska M, Dobosz S, Basiak W. A serological and epidemiological evaluation of risk factors for toxocariasis in children in central Poland. *J Helminthol* 2008; 82: 123-7.
- Zoli AP, Nguekam, Shey-Njila O, *et al.* Neurocysticercosis and epilepsy in Cameroon. *Trans R Soc Trop Med Hyg* 2003; 97: 683-6.