

## RESEARCH NOTE

# BIOEFFICACY OF A LONG-LASTING INSECTICIDE IMPREGNATED NET AGAINST *ANOPHELES MACULATUS* THEOBALD AND *CULEX QUINQUEFASCIATUS* SAY

Tan Swee Beng, Indra Vythilingam and Lee Han Lim

Medical Entomology Unit, WHO Collaborating Centre for Vectors, Institute for Medical Research, Kuala Lumpur, Malaysia

**Abstract.** This study was conducted to evaluate the bioefficacy and effects of unwashed and washed PermaNet® 2.0 nets treated with extrinsic heat during washing against mosquitoes under laboratory condition. The effectiveness of the washed nets was bioassayed soon after washing using the WHO cone test method. The results indicated that the treated nets exhibited high durability to washing (five washes) and persistent bioefficacy against *An. maculatus* on both unwashed and washed nets. The unwashed net exhibited mortality of  $98.33 \pm 2.06\%$ , with KT50 and KT90 ranging from 17.57-to-66.38 minutes and 34.65-to-311.30 minutes, respectively ( $n=9$ ) for >44 months. Nets washed at 30°C for three successive washes exhibited mortality of  $98.00 \pm 2.65\%$  ( $n=3$ ) for 7 months. Twenty-one months later, two washes at 80°C on the same nets caused a mortality of  $87.50 \pm 3.54\%$  ( $n=2$ ). The overall cone mortality of five washes over a total wash period of 33 months was  $94.75 \pm 6.85\%$  with KT50 and KT90 ranging from 23.98 to 58.25 minutes and 51.63 to 230.21 minutes, respectively ( $n=5$ ). However, the treated net exhibited lower bioefficacy against *Culex quinquefasciatus* on the unwashed nets with mortality of 9%. KT50 and KT90 ranged from 129.17 to 136.78 minutes and 315.35 to 716.05 minutes, respectively, over an unwashed period of 3 months ( $n=1$ ), and washed nets had a mortality of 30%; KT50 of 64.04 minutes and KT90 of 347.56 minutes for only one wash at 30°C ( $n=1$ ). Application of extrinsic heat treatment during washing at 30°C followed by increased heat at 80°C on the same net did not enhance any significant increase in mortality of *An. maculatus* and *Culex quinquefasciatus* ( $p>0.05$ ). There was a difference in the delayed knock down time soon after washing at 80°C. Based on these promising results, the PermaNet® 2.0 should be field-tested.

**Keywords:** PermaNet®, bioefficacy, knock down time, laboratory evaluation, WHO cone test, Malaysia

---

Correspondence: Dr Lee Han Lim, Medical Entomology Unit, WHO Collaborating Centre for Vectors, Institute for Medical Research, Jalan Pahang, 50588 Kuala Lumpur, Malaysia.  
Tel/Fax: 6 03 26162688  
E-mail: leehl@imr.gov.my

## INTRODUCTION

Insecticide-treated nets (ITN), especially long-lasting nets (LLN) are widely used in malaria control. Treated nets are useful and provide protection to the people; however, the retreatment of nets

has always posed serious problems in the implementation of ITN. Retreatment rates of nets are very low due to several factors. Many countries do not have the resources for retreatment; these are the cost of the insecticides as well as a lack of understanding of its importance as a form of personal protection proven to reduce severe disease in endemic regions (CDC, 2014). People also do not like to have their nets treated in bulk together with others (Vythilingam *et al*, 2003). Therefore, to avoid the need for periodic re-treatment, it would be advantageous for nets to be able to retain insecticidal efficacy for years and withstand repeated washing (N'Guessan *et al*, 2001).

PermaNet® 2.0 (VestergaardFrandsen, Lausanne, Switzerland) is a type of net pretreated in the factory with deltamethrin, a pyrethroid, which is a neurological insecticide rapidly absorbed by the legs of the mosquito while in contact with the insecticide treated nets. Mosquitoes often dispose off the affected legs when the poison is detected. Therefore, with only three or fewer legs, they can no longer fly, land, and bite, thereby reducing man/vector contact and overall vector density. The World Health Organization (WHO) recommended permethrin as one of the pyrethroids to be widely used in vector control programs (WHO, 1985).

The aim of this study was to evaluate the bioefficacy of the treated net against mosquitoes under laboratory conditions and the efficacy of washed nets with extrinsic heat at 30°C followed by 80°C on the same nets during washing.

## MATERIALS AND METHODS

### Study site

The study was conducted at the main laboratory of the Medical Entomology

Unit/WHO Collaborating Centre for Vectors, Infectious Disease Research Centre, Institute for Medical Research, Kuala Lumpur. The temperature and humidity was maintained at 27±2°C and 80±2%, respectively with a photoperiod of 12:12 hours (L:D).

### Mosquito net tested

PermaNet® 2.0 (VestergaardFrandsen, Lausanne, Switzerland) was made of 75 denier polyester with a mesh size of 25 holes/cm<sup>2</sup> and treated in the factory with 55 mg/m<sup>2</sup> deltamethrin. The size of the net was 160 cm x 150 cm x 180 cm (width x height x length). A similar but untreated net was also tested as a control net. Four samples of the net (25 x 25 cm) were cut from each of the nets and used in the test.

### Mosquitoes tested

The mosquitoes used in the study were laboratory-bred *Anopheles maculatus* Theobald and *Culex quinquefasciatus* Say, which were susceptible to insecticides including pyrethroid. The mosquitoes were fed with 10% sucrose solution and maintained in the insectarium at 27±2°C and 80±2% RH with 12:12 hours (L:D) photoperiod regime.

### Laboratory washing procedure

The net samples (25 cm x 25 cm) were individually introduced into 1 liter beakers containing 0.5 liter distilled water, with 2 g/l soap (Savon de Marseille, pH 10-11) fully dissolved. The beakers were then placed into a shaking water bath at 30°C and 80°C, and shaken for 10 minutes at 155 rotations per minute. The net samples were then removed and rinsed twice for 10 minutes in clean distilled water in the same shaking condition as stated above. After washing and drying at room temperature, the net samples were tested to determine effectiveness soon after washing. In this trial, the same nets were

washed three times at 30°C followed by two final washes at 80°C and shaken for 10 minutes at 155 rotations per minute. The purpose of washing the net in hot water was to determine if heat could regenerate the activity of the net after washing.

#### Bioassay of mosquitoes

Four pieces of net were cut from each net. The standard WHO plastic cones were placed on each net and labeled 1-4, accordingly. Five sugar-fed, 2-to-5 days old female mosquitoes were introduced into each cone. The cones were then plugged with cotton wool and covered with a black cloth to ensure total contact of the mosquitoes with the treated net. The exposure time was 3 minutes, after which the mosquitoes were transferred into clean paper cups accordingly, and mortality rate was recorded at 10-minute intervals until 60 minutes.

The mosquitoes were held in these paper cups for 24 hours at 27±2°C and 80±2% RH with 12:12 hours (L:D) photoperiod, and 10% sucrose solution was provided. Each sample was tested using a total of 25 mosquitoes, and four such replications were carried out. Tests were also conducted on the two pieces of control (untreated) nets in parallel to the test and delayed mortality observed after 24 hours. Test net materials used for residual were kept indoor in the laboratory at ambient temperature.

#### Data analysis

The number of dead mosquitoes was recorded. If the control mortality was between 5%-20%, the mortalities will be corrected by Abbott's formula (Abbott, 1925):

$$\frac{\% \text{treated mortality} - \% \text{control mortality}}{100\% - \% \text{control mortality}} \times 100\%$$

When results obtained in test where control mortalities exceed 20%, the test

would be discarded and the experiment repeated. All data were expressed as the mean ± SEM. A value of  $p < 0.05$  was considered significant. Data obtained from the test were subjected to statistical analysis using SPSS® (version 16.0, SPSS: Chicago, IL), while knockdown data were subjected to probit analysis using similar software to obtain the KT50 and KT90 values.

## RESULTS

#### *An. maculatus*

With unwashed treated nets, *An. maculatus* exhibited KT50 and KT90 ranging from 17.57-to-66.38 minutes and 34.65-to-311.30 minutes, respectively for >44 months (Table 1). Bioefficacy of knock down time of washed treated net against *An. maculatus* exhibited KT50 and KT90 ranging from 23.98-to-58.25 minutes and 51.63-to-230.21 minutes, respectively for five washes over a test period of 33 months with a difference in the delayed knock down time of 34.27 minutes soon after washing at 80°C (Table 2).

The residual activity of treated net against *An. maculatus* on unwashed nets was >44 months with a mean mortality of 98.33±2.06% ( $n=9$ ), while the nets washed at 30°C for three successive washes exhibited mortality of 98.00±2.65% ( $n=3$ ) for 7 months and 21 months later (from 28<sup>th</sup>-to 33<sup>rd</sup> month). Two washes at 80°C on the same nets exhibited mortality of 87.50±3.54% ( $n=2$ ). However, the overall mortality of five washes (30°C followed by 80°C) was 94.75±6.85% ( $n=5$ ) over a total wash period of 33 months (Table 4). There was no significant difference between unwashed and washed treated nets ( $p > 0.05$ ).

#### *Culex quinquefasciatus*

*Culex quinquefasciatus* exhibited lower

Table 1  
Knockdown rate of *An. maculatus* on unwashed nets.

Unwashed (Months)	KT50 (Min) (95% Confidence limit)	KT90 (Min) (95% Confidence limit)	Regression line
0	17.57 (14.37-20.55)	34.65 (29.21-44.39)	$y=4.35 x-5.41$
3	21.01 (19.21-22.75)	45.61 (41.44-51.24)	$y=3.81 x-5.04$
8	25.39 (15.37-34.80)	71.95 (48.33-233.45)	$y=2.83 x-3.98$
16	14.40 (12.35-16.28)	41.13 (36.38-47.97)	$y=2.81 x-3.26$
29	46.38 (34.56-84.87)	163.50 (87.78-1,683.85)	$y=2.34 x-3.90$
34	52.83 (43.80-71.11)	433.93 (231.61-1,392.23)	$y=1.40 x-2.41$
39	41.69 (35.62-50.97)	280.85 (172.41-650.04)	$y=1.55 x-2.51$
41	54.08 (41.08-114.39)	150.61 (84.76-1,769.17)	$y=2.88 x-4.99$
>44	66.38 (44.99-336.56)	311.30 (121.83-56,467.30)	$y=1.91 x-3.48$

Table 2  
Knockdown rate of *An. maculatus* on washed nets.

Washed (Months)	KT50 (Min) (95% Confidence limit)	KT90 (Min) (95% Confidence limit)	Regression line
1 <sup>st</sup> 30°C	23.98 (22.07-25.84)	51.63 (46.82-58.24)	$y=3.85 x 5.31$
2 <sup>nd</sup> 30°C	23.61 (14.22-31.73)	59.35 (41.98-151.26)	$y=3.20 x-4.40$
3 <sup>rd</sup> 30°C	25.42 (23.25-27.56)	61.13 (54.38-70.96)	$y=3.36 x-4.73$
4 <sup>th</sup> 80°C	59.15 (50.71-73.78)	250.51 (167.31-483.01)	$y=2.04 x-3.62$
5 <sup>th</sup> 80°C	58.25 (45.04-100.12)	230.21 (122.63-1,272.99)	$y=2.15 x-3.79$

Table 3  
Knockdown rate of *Cx. quinquefasciatus* on unwashed and washed nets.

Unwashed (Months)	KT50 (Min) (95% Confidence limit)	KT90 (Min) (95% Confidence limit)	Regression line
0	136.78 (94.79-285.37)	716.05 (327.34-3,635.30)	$y=1.78 x-3.81$
3	129.17 (91.35-338.61)	315.35 (168.05-1,895.00)	$y=3.31 x-6.98$
1 wash, 30°C	64.04 (53.08-85.47)	347.56 (207.95-844.21)	$y=1.75 x-3.15$

bioefficacy of knock down times of KT50 and KT90 ranging from 129.17-to-136.78 minutes and 315.35-to-716.05 minutes over an unwashed period of 3 months. Bioefficacy of washed treated nets exhibi-

ted knock down times of KT50 (64.04 minutes) and KT90 (347.56 minutes) for only one wash at 30°C with a difference in the knock down time of 65.13 minutes soon after washing (Table 3).

Table 4  
Mortality of *An. maculatus* and *Cx. quinquefasciatus* on unwashed and washed nets.

Treatment	Months	Species	Replicate (n)	Mortality (%)	p-value
Unwashed	>44	<i>An. maculatus</i>	9	98.33 ± 2.06	0.00
3 washes, 30°C	7	<i>An. maculatus</i>	3	98.00 ± 2.65	0.00
2 washes, 80°C	28 <sup>th</sup> - 33 <sup>rd</sup>	<i>An. maculatus</i>	2	87.50 ± 3.54	0.0018
Overall washed 30°C and 80°C	33	<i>An. maculatus</i>	5	94.75 ± 6.85	0.00
Unwashed/washed		<i>An. maculatus</i>	5	3.00 ± 3.94	0.164
Unwashed	1	<i>Cx. quinquefasciatus</i>	1	9.00	nil
1 wash, 30°C	1	<i>Cx. quinquefasciatus</i>	1	30.00	nil

The mortality of *Cx. quinquefasciatus* on unwashed nets was 9% ( $n=1$ ), while the nets washed at 30°C exhibited mortality of 30% ( $n=1$ ) (Table 4). Because the mortality was so low, no further test was carried out at 80°C.

#### DISCUSSION

The bioefficacy of insecticide treated bednets in preventing morbidity and mortality from malaria endemic countries had been studied and reported by several researchers (Alonso *et al*, 1993; Dolan *et al*, 1993; Jaenson *et al*, 1994; Stich *et al*, 1994; D'Alessandro *et al*, 1995a, 1995b; Binka *et al*, 1996; Nevillet *et al*, 1996; Rowland *et al*, 1996; Goodman *et al*, 1999; Armstrong-Schellenberg *et al*, 2001; Guyatt and Snow, 2002; Lengeler, 2004; Rafinejad *et al*, 2008).

Our cone bioassay indicated that PermaNet exhibited high durability to washing (five washes) and persistent bioefficacy against *An. maculatus* on both unwashed, with mortality of 98.33±2.06% ( $n=9$ ) for >44 months, and nets washed at 30°C for three successive washes with mortality of 98.00±2.65% ( $n=3$ ) for 7 months and 21 months later (from 28<sup>th</sup>-to-33<sup>rd</sup>-months), followed by two washes at

80°C on the same nets with a mortality of 87.50±3.54% ( $n=2$ ) over a washed period of 33 months.

However, the overall cone mortality of five washes over a total washed period of 33 months was 94.75±6.85% ( $n=5$ ). Our study concurred with previous research done in Colombia and Bolivia using a cone test for the evaluation of PermaNet<sup>®</sup>, lambda-cyhalothrin, deltamethrin, and alpha cypermethrin impregnated polyester nets against *Anopheles* sp where 100% mortality was reported after three washes (Gonzalez *et al*, 2002).

However, PermaNet<sup>®</sup> 2.0 exhibited lower bioefficacy against *Cx. quinquefasciatus* on unwashed and washed nets. Our study concurred with Hougard *et al* (2003) that *Cx. quinquefasciatus* exhibited lower susceptibility to most chemical insecticides.

In comparison to the tunnel test, higher mortality was observed in cone bioassay that has forced tarsal contact for mosquitoes to rest on the insecticide treated nets. Our previous study (Tan *et al*, 2014) on tunnel mortality using the same nets concurred with the results of Bhatt *et al* (2005) and Rafinerad *et al* (2008) where higher mortality of their cone and




Fig 1–WHO cone test on bioefficacy of unwashed and washed PermaNet® 2.0 against *An. maculatus* and *Cx. quinquefasciatus*.

tube test on six times washed nets were observed and the percentage of 1 hour knockdown on unwashed and six times washed nets was 100% in both cone and tubes. In tunnel tests, 44% and 74% mortality rates were recorded on unwashed and six times washed nets, respectively. Our tunnel mortality on the same nets exhibited 74.36±25.77% (n=14) mortality on unwashed and 75.00±18.46% (n=10) after five washes. Cone mortality exhibited higher mortality of 98.33±2.06% (n=9) (unwashed) and 94.75±6.85% (n=5) (five washes). The variation in the magnitude of decline in tunnel mortality may be due to the large space of tunnel; therefore, mosquitoes are not obliged to come into contact with the nets impregnated with insecticides (Rafinejad *et al*, 2008).

This study also suggested that there was no significant difference in knockdown time and mortality of *An. maculatus* and *Cx. quinquefasciatus* between unwashed and washed nets ( $p>0.05$ ). Therefore, applica-

tion of extrinsic heat treatment of 30°C followed by 80°C on the same nets during washing did not enhance any significant decrease in the knock down time and increase in mortality of both *An. maculatus* and *Cx. quinquefasciatus*.

Since washing at 30°C was done only once, and no further washing was conducted at 80°C, future study on larger sample size of *Cx. quinquefasciatus* should be conducted to obtain better and reliable results.

Lindblade *et al* (2005) concluded that PermaNet® 2.0 performed significantly better than other conventional nets. The wash resistance offered by PermaNet® 2.0 was much better and long lasting (Rafinejad *et al*, 2008). Graham *et al* (2005) found 97% mortality against *Anopheles* in Pakistan, Iran, and Tanzania trials using PermaNet® 2.0 after 21 washes, whereas our study found that after 5 washes, PermaNet® 2.0 still remained effective in killing susceptible lab-strain *An. maculatus*

of up to 94.75±6.85% ( $n=5$ ). Based on data obtained, we suggest PermaNet® 2.0 be field-tested to assess its impact on malaria.

#### ACKNOWLEDGEMENTS

The authors would like to thank the Director General of Health, Malaysia and Dr Shahnaz Bt Murad, Director, Institute for Medical Research, Kuala Lumpur for their support. Thanks are also due to Mr AA Abdul Malek, Mr S Subramaniam, and Mr T Mustami for their kind assistance.

#### REFERENCES

- Abbott WS. A method of computing the effective of an insecticide. *J Econ Entomol* 1925; 18: 265-7.
- Alonso PL, Lindsay SW, Armstrong-Schellenberg JRM, *et al.* A malaria control trial using insecticide-treated bednets and targeted chemoprophylaxis in a rural area of the Gambia, West Africa. *Trans R Soc Trop Med Hyg* 1993; 87 (suppl): 37-44.
- Armstrong-Schellenberg JRM, Abdulla S, Nathan R, Mukasa O, Marchant TJ. Effect of large scale social marketing of insecticide-treated nets on child survival in rural Tanzania. *Lancet* 2001; 357: 1241-7.
- Bhatt RM, Yadav RS, Adak T, Babu CJ. Persistence and wash-resistance of insecticidal efficacy of nettings treated with deltamethrin tablet formulation (K-O TAB) against malaria vectors. *J Am Mosq Control Assoc* 2005; 21: 54-8.
- Binka FN, Kubaje A, Adjuik M, *et al.* Impact of permethrin impregnated bednets on child mortality in Kassena Nankana District, Ghana: a randomized control trial. *Trop Med Int Health* 1996; 1: 147-54.
- Centers for Disease Control and Prevention (CDC). 2014 insecticide-treated bed nets. Atlanta: CDC, 2014. [Cited 2014 Mar 27]. Available from: [http://www.cdc.gov/malaria\\_worldwid/reduction/itn.html](http://www.cdc.gov/malaria_worldwid/reduction/itn.html)
- D'Alessandro U, Olaleye BO, McGuire W, *et al.* Mortality and morbidity from malaria in Gambian children after introduction of impregnated bednet programme. *Lancet* 1995a; 345: 479-83.
- D'Alessandro U, Leach A, Drakeley CJ, *et al.* A comparison of the efficacy of insecticide-treated and untreated bed nets in preventing malaria in Gambian children. *Trans R Soc Trop Med Hyg* 1995b; 89: 596-8.
- Dolan G, TerKuile FO, Jacoutot V, *et al.* Bednets for prevention of malaria and anaemia in pregnancy. *Trans R Soc Trop Med Hyg* 1993; 87: 620-6.
- González J, Kroeger A, Aviña AI, Pabón E. Wash resistance of insecticide-treated materials. *Trans R Soc Trop Med Hyg* 2002; 96: 370-5.
- Goodman CA, Coleman PG, Mills AJ. Cost-effectiveness of malaria control in sub-Saharan Africa. *Lancet* 1999; 354: 378-85.
- Graham K, Kayedi MH, Maxwell C, *et al.* Multi-country field trials comparing wash-resistance of PermaNet and conventional insecticide-treated nets against anopheline and culicine mosquitoes. *Med Vet Entomol* 2005; 19: 72-83.
- Guyatt HL, Snow RW. The cost of not treating bednets. *Trends Parasitol* 2002; 18: 12-6.
- Hougard JM, Duchon S, Darriet F, Zaim M, Rogier C, Guillet P. Comparative performances, under laboratory conditions, of seven pyrethroid insecticides used for impregnation of mosquito nets. *Bull World Health Organ* 2003; 81: 324-33.
- Jaenson TGT, Gomes MJ, Barreto dos Santos RC, Retranca V, Fortini D, Crato J. Control of endophagic *Anopheles* mosquitoes and human malaria in Guinea Bissau, West Africa by permethrin treated bednets. *Trans R Soc Trop Med Hyg* 1994; 88: 620-4.
- Lindblade KA, Dotson E, Hawley WA, *et al.* Evaluation of long-lasting insecticidal nets after 2 years of household use. *Trop Med Int Health* 2005; 11: 1141-50.
- Lengeler C. Insecticide-treated bednets and curtains for preventing malaria. *Cochrane Database Syst Rev* 2004; (2): CD000363.
- Nevill CG, Some ES, Mung'ala VO, *et al.* Insec-

- ticide-treated bednets reduce mortality and severe morbidity from malaria among children on the Kenyan coast. *Trop Med Int Health* 1996; 1: 139-46.
- N'Guessan R, Darriet F, Doanniom JM, Chandre F, Carnevale P. Olyset-Net efficacy against pyrethroid-resistant *Anopheles gambiae* and *Culex quinquefasciatus* after 3 years' field use in Cote d'Ivoire. *Med Vet Ent* 2001; 15: 97-104.
- Rafinejad J, Vatandoost H, Nikpoor F, et al. Effect of washing on the bioefficacy of insecticide-treated nets (ITNs) and long lasting insecticidal nets (LLINs) against main malaria vector *Anopheles stephensi* by three bioassay methods. *J Vector Borne Dis* 2008; 45: 143-50.
- Rowland M, Bouma M, Ducornez D, et al. Pyrethroid-impregnated bednets for personal protection against malaria for Afghan refugees. *Trans R Soc Trop Med Hyg* 1996; 90: 357-61.
- Stich AH, Maxwell CA, Haji A, et al. Insecticide-impregnated bednets reduce malaria transmission in rural Zanzibar. *Trans R Soc Trop Med Hyg* 1994; 88: 150-4.
- Tan SB, Vythlingam I, Lee HL. Bioefficacy of a long-lasting insecticide impregnated net on blood feeding inhibition of *Anopheles maculatus* Theobald and *Culex quinquefasciatus* Say. *Southeast Asian J Trop Med Public Health* 2014; 45: 598-609.
- Vythilingam I, Sutjipto SKM, Tan SB. The effectiveness of kits for home dipping for treatment of mosquito nets. *Trop Biomed* 2003; 20: 19-25.
- World Health Organization (WHO). Safe use of pesticides. *WHO Techn Rep Ser* 1985; 720.